

Cohomology of Projective Spaces

Feng Ling

The University of Texas At Austin

FLing@utexas.edu

April 26, 2015

Derived Functors

Knowing sheaf cohomology of things is very useful in solving problems in multi-variable complex analysis, classical and abstract algebraic geometry

We can define the sheaf cohomology in generality using the concept of derived functors.

Definition

Given an left exact functor $F : \mathfrak{A} \rightarrow \mathfrak{B}$ and some injective resolution I^\cdot of X in \mathfrak{A} with induced morphisms $d^i : F(I^i) \rightarrow F(I^{i+1})$, a right derived functor $R^i F$ is defined to be $\ker d^i / \text{im } d^{i-1}$.

Of course this definition is also valid for left derived functors and projective resolutions etc.

Abelian Group Sheaf Cohomology and Mayer-Vietoris

Applying the above to the left exact global section functor $\Gamma : \mathfrak{Ab}(X) \rightarrow \mathfrak{Ab}$ of a topological space X , we can get our corresponding cohomology of sheaves of abelian groups $H^i(X, \mathcal{F}) = R^i\Gamma$ from injective resolution I^\bullet of sheaf \mathcal{F} on X .

Computing cohomologies directly from an explicit injective resolution for every (X, \mathcal{F}) is usually difficult. But we can define the so-called Čech cohomology to help us. To motivate its definition, let's first consider the Mayer-Vietoris Sequences.

Theorem (Mayer-Vietoris Sequences)

If $X = U \cup V$, \mathcal{F} a sheaf on X , and $i : U \rightarrow X$, $j : V \rightarrow X$, $k : U \cap V \rightarrow X$ are the inclusion maps, we get a long exact sequence of cohomologies $0 \rightarrow H^0(X, \mathcal{F}) \rightarrow H^0(U, \mathcal{F}) \oplus H^0(V, \mathcal{F}) \rightarrow H^0(U \cap V, \mathcal{F}) \rightarrow H^1(X, \mathcal{F}) \rightarrow \dots$

Čech Cohomology

Likewise, we define the Čech cohomology of (X, \mathcal{F}) by using $\mathfrak{U} = \{U_i\}_{i \in I}$ an ordered open cover of X .

The cohomologies would be based on the cochain of complexes

$$C^p := \prod_{i_0 < \dots < i_p} \mathcal{F}(U_{i_0} \cap \dots \cap U_{i_p})$$

which forms a sequence $0 \rightarrow C^0 \rightarrow C^1 \rightarrow \dots \rightarrow C^p \rightarrow \dots$ with coboundary map $d^p : C^p \rightarrow C^{p+1}$ defined as

$$d^p(\dots, \alpha_{i_0, \dots, i_p}, \dots)_{i_0, \dots, i_{p+1}} := \sum_{k=0}^{p+1} (-1)^k \alpha_{i_0, \dots, \hat{i}_k, \dots, i_p} \Big|_{U_{i_0} \cap \dots \cap U_{i_{p+1}}}$$

Cohomology of Projective Spaces

For projective spaces, we have $X = \mathbb{P}_A^r = \text{Proj } S$ where $S = A[x_0, \dots, x_r]$ and a quasi-coherent sheaf $\mathcal{F} = \bigoplus_{n \in \mathbb{Z}} \mathcal{O}(n)$ where $\mathcal{O}(n) = \widetilde{S}(n)$.

This structure naturally admits an (affine) open covering $\mathcal{U} = \{\text{Spec } S_{(x_i)}\}_{0 \leq i \leq r}$ where $S_{(x_i)} = A[\frac{x_0}{x_i}, \dots, 1, \dots, \frac{x_r}{x_i}]$ is the localized degree 0 coordinate ring. Now we have the Čech complexes

$$\begin{aligned} C^p &= \prod_{i_0 < \dots < i_p} \mathcal{F}(\text{Spec } S_{(x_{i_0} \dots x_{i_p})}) \\ &= \prod_{i_0 < \dots < i_p} \bigoplus_{n \in \mathbb{Z}} \mathcal{O}(n)(\text{Spec } S_{(x_{i_0} \dots x_{i_p})}) \\ &= \bigoplus_{n \in \mathbb{Z}} \prod_{i_0 < \dots < i_p} \widetilde{S_{(x_{i_0} \dots x_{i_p})}}(n) \\ &\cong \prod_{i_0 < \dots < i_p} S_{x_{i_0} \dots x_{i_p}} \end{aligned}$$

Computing the Čech cohomology

So we have the sequence

$$0 \rightarrow \prod_i S_{x_i} \xrightarrow{d^0} \prod_{i,j} S_{x_i, x_j} \xrightarrow{d^1} \cdots \xrightarrow{d^{r-2}} \prod_i S_{x_0, \dots, \hat{x}_i, \dots, x_r} \xrightarrow{d^{r-1}} S_{x_0, \dots, x_r} \rightarrow 0$$

which induces the Čech cohomology sequence.

$$0 \rightarrow \check{H}^0(X, \mathcal{F}) = \ker d^0 \rightarrow \check{H}^1(X, \mathcal{F}) \rightarrow \cdots \rightarrow \check{H}^i(X, \mathcal{F}) = \frac{\ker d^i}{\text{im } d^{i-1}} \rightarrow \cdots \rightarrow \check{H}^{r-1}(X, \mathcal{F}) \rightarrow \check{H}^r(X, \mathcal{F}) = \text{coker } d^{r-1} \rightarrow 0$$

with which we can start our calculation...

Useful Facts

Theorem (1)

If X is affine and \mathcal{F} a quasi-coherent sheaf, then $H^p(X, \mathcal{F}) = 0 \forall p > 0$.

Theorem (2)

If X is noetherian separated scheme, \mathfrak{U} an open affine cover and \mathcal{F} a quasi-coherent sheaf, then $\check{H}^p(\mathfrak{U}, \mathcal{F}) \rightarrow H^p(X, \mathcal{F})$ are isomorphisms for $p \geq 0$.

Theorem (3)

For $Y \hookrightarrow X$ a closed subscheme with inclusion map i , we have the exact sequence $0 \rightarrow \mathcal{I}_Y \rightarrow \mathcal{O}_X \rightarrow i_\mathcal{O}_Y \rightarrow 0$*

Theorem (4)

If a module M have an isomorphism to itself by multiplication of f , then $M_f = 0$ implies $M = 0$.

Table of $H^p(\mathbb{P}_A^r, \mathcal{O}(n))$

n	$p = 0$	$p = 1$...	$p = r - 1$	$p = r$
m	$A(\dots, f_i, \dots)$	0	...	0	0
1	$A(x_0, \dots, x_r)$	0	...	0	0
0	A	0	...	0	0
-1	0	0	...	0	0
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots
$-r$	0	0	...	0	0
$-r - 1$	0	0	...	0	B
$-r - 2$	0	0	...	0	$B(x_0^{-1}, \dots, x_r^{-1})$
$-r - 1 - m$	0	0	...	0	$B(\dots, f_i^{-1}, \dots)$

where $B = A\left(\frac{1}{(x_0 \cdots x_r)}\right)$, $m > 1$, and f_i are the degree m monomials in x_0, \dots, x_r .



Robin Hartshorne (1977)

Algebraic Geometry

And most importantly my mentor: Zhu, Yuecheng!